

AMPROL 12% SOLUTION

(Amprolium)
Treatment and prevention of coccidiosis in poultry
1, 5 Litres Can containing an oral 12% Amprolium Hydrochloride

COMPOSITION

100ml of the oral solution contain:

Amprolium Hydrochloride 12.0g (corresponding to 10.6g of Amprolium)

Excipients qs 100ml

THERAPEUTIC INDICATIONS

Treatment and prevention of coccidiosis in poultry due to infections of *Eimeria spp* sensitive to amprolium.

ROUTE AND METHOD OF ADMINISTRATION

Oral route. Administer via the drinking water.

Dosage:

Treatment:

For usual outbreak, 1ml of Amprolium 12% Solution per litre of drinking water for 5 to 7 days.

For sever outbreak, 2ml of Amprolium 12% Solution per litre of drinking water for 5 to 7 days.

Maintenance treatment:

0.5ml of Amprolium 12% Solution per litre of drinking water for 1 to 2 weeks.

Prophylaxis:

0.5ml of Amprolium 12% Solution per litre of drinking water for the duration of risk periods.

PRECAUTIONS FOR USE

- Shake before use
- Do not smoke or eat while handling the product
- Wash hands after handling the product
- Keep out of reach of children

WITHDRAWAL PERIOD

Nil.

PRESENTATION

1, 5 Litres Can.

Manufactured By:



23 Rue du Prieuré, Saint-Herblon
44150 Vair-sur-Loire, France